REVIEW ARTICLE

A Comprehensive Review on Phytochemical Profile of *Acacia auriculiformis*

Ruchi Dobariya^{1*}, Vaibhavi Savalia¹, Krishna Raninga², Pravin Tirgar¹, Jigar Savaliya³, Anjali Parmar²

DOI: 10.18811/ijpen.v10i03.04

Abstract

Acacia auriculiformis A. Cunn. Ex Benth is a globally distributed perennial shrub known for its wide-ranging medicinal properties and therapeutic uses, with a low toxicity profile. Traditionally, it has been used to treat conditions such as rashes, rheumatism, allergies, eye irritation, and pain. Its pharmacological activities include strong antimicrobial and antioxidant effects, which help combat infections and reduce oxidative stress. The plant has shown promising antimalarial and anti-filarial activities, making it useful for treating parasitic diseases like malaria and lymphatic filariasis. Additionally, A. auriculiformis exhibits cytocidal and antimutagenic properties, suggesting potential for cancer prevention and treatment. It also has spermicidal, wound healing, and hepatoprotective effects, further broadening its therapeutic applications. Of particular interest is its antidiabetic activity, demonstrating potential in managing blood sugar levels and improving insulin sensitivity. These pharmacological effects are attributed to the plant's bioactive compounds, such as flavonoids and tannins. This review highlights the plant's phytochemical composition, pharmacogenetic factors, and its broad spectrum of therapeutic activities, making A. auriculiformis a valuable resource for both traditional and modern medicine. Ongoing research is expected to further explore its potential for treating a variety of health conditions.

Highlights:

- The medicinal applications are broad, and the plant is used for treating irritation, rashes, rheumatism, allergies, and pain relief.
- It has extensive pharmacological effects, including antimicrobial, antioxidant, antimalarial, and antidiabetic properties.
- Its phytochemical profile is rich and contains bioactive compounds with significant therapeutic potential.
- The pharmacogenetic insights explore genetic influences on the plant's efficacy and safety.
- The plant demonstrates low toxicity and high effectiveness, with minimal adverse effects while providing therapeutic benefits.

Keywords: Acacia auriculiformis, Pharmacology, Phytochemicals, Nutritional value.

ISSN: 2454-1117 (Print), 2455-202X (Online)

INTRODUCTION

The Mimosaceae family includes the rapidly spreading Acacia auriculiformis A. Cunn, a deciduous or evergreen tree that may grow to a height of 30 meters. It has high concentrations of methyl glucuronic acid, glucuronic acid, galactose, arabinose, and rhamnose. Because of the tannins and triterpenoid saponins present, it is said to have spermicidal, filaricidal, and central nervous system depressant properties (Srivastava and Singh, 2024; Prayogo *et al.*, 2024; Thorat and Kadre, 2024).

International Journal of Plant and Environment (2024);

The Fabaceae family includes Acacia auriculiformis (A. auriculiformis) A.Cunn. ex Benth., a medium-sized, straight, deciduous, or evergreen tree that may grow to a height of up to 30 meters. It is typically found in India's parks and roadside areas. Acacia's common name comes from the Greek word "akis," which means "spike" or "point." In contrast, the latin word "auricula" denotes a creature's outer ear and is a specific name, whereas the word "forma" denotes a frame, figure, or shape. The tree is originally from Australia and was brought to West Bengal, India, in 1946.

The primary aim of the study was to showcase the most recent pharmacological and phytochemical studies conducted on the *A. auriculiformis* plant to date. Using the keywords "*A. auriculiformis* phytochemistry," "*A. auriculiformis* pharmacology," and "*A. auriculiformis* patents" in the SciFinder, PubMed, Scopus, and Google Scholar databases, a comprehensive and relevant literature search was performed. Results from the literature

¹School of Pharmacy RK University Rajkot., Dhananjay, Gokul nagar Talala, Gir Somnath, Gujarat, India.

²Parul Institute of Pharmacy, Limda Waghodia, Gujarat, India.

³Pandit Dindayal Medical College, Rajkot, Gujarat, India.

***Corresponding author:** Ruchi Dobariya, School of Pharmacy RK University Rajkot., Dhananjay, Gokul nagar Talala, Gir Somnath, Gujarat, India, Email: ruchidobariya4499@gmail.com

How to cite this article: Dobariya, R., Savalia, V., Raninga, K., Tirgar, P., Savaliya, J., Parmar, A. (2024). A Comprehensive Review on Phytochemical Profile of *Acacia auriculiformis*. International Journal of Plant and Environment. 10(3), 36-47.

Submitted: 02/07/2024 Accepted: 07/10/2024 Published: 30/11/2024

search showed papers published. Following that, specific papers on pharmacology and phytochemistry were carefully screened, chosen, and examined without regard to the chronological order to compile the current manuscript (Ge *et al.*, 2015).

The current review delves into traditional, ethnobotanical, phytochemical, pharmacological, and nutraceutical knowledge. Modern pharmacological research and the presence of several important phytochemicals back up ethnobotanical and traditional assertions. The presence of essential proteins, vitamins, amino acids, and minerals demonstrates *A. auriculiformis* importance as a nutraceutical.

MATERIALS AND METHOD

Comprehensive information related to *A. auriculiformis* was collected using the keywords, botanical name of *A. auriculiformis* and common name of *A. auriculiformis* in various electronic databases like Pubmed, google Scholar, and Science Direct. Additionally, some books were also referenced.

Taxonomy

Auriculiformis is formally referred to as *A. auriculiformis A.Cunn. ex Benth.* Additional informal designations for this plant encompass papuan wattle, dal moth, earleaf acacia, auri, earpod wattle, black wattle, tan wattle, northern black wattle, and darwin black wattle. In telugu, it is known as minnumaan, kondamanu, seema babul, and maha babul; in tamil as kaththi karuvel; in kannada as aurculis; in marathi as akashia; and in Australian babool.

A. auriculiformis A. Cunn. Ex Benth, belonging to the family fabaceae (Leguminosae), is a prominent member of the plant kingdom. Classified under the domain Eukaryota and the kingdom Plantae, it falls within the subkingdom Viridiplantae and the infrakingdom streptophyta. This species is part of the phylum Spermatophyta, specifically the subphylum Angiospermae, and belongs to the class Magnoliopsida (Dicotyledonae). Within the subclass Rosidae and Superorder Rosanae, A. auriculiformis is situated in the order Fabales. It is categorized under the division Tracheophyte and subdivision Spermatophytina, and its genus is Acacia Mill.

A. auriculiformis Cunn.ex Benth., orth. Var., A. moniliformis Griseb., and Racosperma auriculiformae (A.Cunn. ex Benth.), pedley are among the synonyms of A. auriculiformis.

Botanical Description

The duration of the season for flowers and fruiting for *A. auriculiformis* differs by region; in Australia, it spans from April to July. Seeds are ripe four to five months later, between August and October. Malaysia experiences floral blooms from February to may, and the harvesting season for mature produce seeds is from October to April. Java, Indonesia, experiences its flowering season from March to June, while Thailand observes the mature seed collection period from August to February. In India, *A. auriculiformis* blooms from December to January the year before and bears fruit from February to March. On the contrary, fruiting and flowering transpire in diverse regions of India between March and December, exhibiting a peak profusion during September and October. Bark, leaves, and fruits of *auriculiformis*—pods comprising seeds and funicles—are utilized in a wide range of biological processes.

In its natural habitat, *A. auriculiformis* attains a maximum height of 25 to 35 m at favorable locations, where a straight bole composes the majority of the tree's stature. Typically 8 to 20 meters in height, it is sporadically a 3 to 5 m tall shrub with a short bole that is densely branched. The bark of juvenile trees is smooth and gray or brown, with intermittent blackening at the base; as the tree matures, it becomes longitudinally fissured and rugged. Phallic-shaped, glabrous, greyish-green, and faintly textured phyllodes are observed. They lengthen to 8 to 20 cm and widen to 1.0 to 4.5 cm. In addition to minutes, densely packed secondary veins, and a specific duct located at the base of the phyllode, three prominent longitudinal veins are evident. These veins run in a parallel line towards the lower margin or in the central region, near the base. The inflorescence, which grows in pairs in the upper axils, is an axillary spike that is interrupted and can reach a maximum length of 8.5 cm.

The flowers are sessile, minute, fragrant, tubular, fivemerous, and pale golden in hue. Numerous, approximately 0.3 cm long filaments adorn the corolla, which can reach a maximum length of 0.2 cm. The ovary is densely pubescent. Approximately, 0.1 cm in length, the calyx is succinctly lobed. The pods have an approximate length of 6.5 cm and a width of 1.5 cm. They are flat, flexible yet unyielding, closely curled into an open coil, glaucous, resinous to a certain degree, and veined transversely with undulating boundaries. They transform from their original straight or curved state to one of random coiling and twisting as they mature. The lustrous black seeds, which measure 0.4 to 0.6 cm in length and 0.3–0.4 cm in width, are arranged transversely within the pod. Each has an elongated, elliptical, or red, yellow, or orange funicle; the areole is substantial and nearly encased.

Botanical Features

In optimal conditions within its natural habitat, *A. auriculiformis* develops into a 25 to 35 m tall tree, with a straight bole encompassing the majority of the tree's height. It is usually an 8 to 20 m tree, although it is sometimes occasionally a 3 to 5 m shrub with a low bole and many branches. In young trees, the bark is smooth and gray or brown, with occasional blackening around the base; as the tree ages, it becomes rough and longitudinally fissured.

Inflorescences, Flowers and Fruits

The inflorescence, which grows in pairs in the upper axils, is an axillary spike that is interrupted and can reach a maximum length of 8.5 cm. The flowers are sessile, minute, fragrant, tubular, five-merous, and pale golden in hue. Numerous approximately 0.3 cm-long filaments adorn the corolla, which can reach a maximum length of 0.2 cm. The ovary is densely pubescent. Approximately 0.1 cm in length, the calyx is succinctly lobed. The pods are approximately measuring 6.5 cm in length and 1.5 cm in breadth. They are flat, flexible yet unyielding, closely curled into an open coil, glaucous, resinous to a certain degree, and veined transversely with undulating boundaries. They transform from their original straight or curved state to one of random coiling and twisting as they mature. The lustrous black seeds, which measure 0.4 to 0.6 cm in length and 0.3 to 0.4 cm in width, are arranged transversely within the pod. Each has an elongated, elliptical, or red, yellow, or orange funicle; the areole is substantial and nearly encased.

Leaves

The phyllodes are falcate, glabrous, greyish-green, and lightly textured. They lengthen to 8 to 20 cm and widen to 1.0 to 4.5 cm. In addition to several minutes, densely packed secondary veins, and a discernible duct located beneath the phyllode, three prominent longitudinal veins are apparent and run in a parallel direction towards the lower margin or in the central region, close to the base. (Ahmadu *et al.*, 2024)

Habitat and Ecology

A. auriculiformis is indigenous to Papua New Guinea, Indonesia and Australia, where it often lives on land. It is found in every state in India except for Arunachal Pradesh, Sikkim, and Jammu and Kashmir. This is due to its quick growth, drought resistance, seasonal tolerance of wet soils, and capacity to grow in poor soils.

Phytochemical Description

This section discusses the phytochemicals isolated from *A*. *auriculiformis* (Tables 1 and 2).

Flavonoids

Several flavonoids have been documented in extracts of *A. auriculiformis*. In the early 1960s, a unique flavan-3,4-diol was extracted from the heartwood of *A. auriculiformis* using paper iontophoresis (Tiho *et al.*, 2024). Additionally, a novel flavan glucoside known as auriculoside (I, Glc= β -D-glucopyranosyl) or 7,3',5'-trihydroxy-4'-methoxyflavan 3'-glucoside, was isolated and demonstrated 80 % CNS depressive activity (Singh, 2024; Ogunniyi *et al.*, 2023). The bark of *A. auriculiformis* contains quercetin [2-(3,4-dihydroxyphenyl)-3,5,7-trihydroxy-4*H*-chromen-4-one] and epicatechin [(2R,3R)-2-(3,4-

		Tuble 1. Showing phytoenemicals isolated nonini. adheano	initial that of statutes	
S. No.	Part used	Extract/Phytoconstituent	Dose tested	Reference
1	Heartwood	lsolated compounds (3,4',7,8-tetrahydroxyflavanone, 4',7,8-trihydroxyflavanone, Teracacidin), methanol, diethyl ether, ethyl acetate, n-butanol extracts.	0.1, 1.0, and 10.0 mg/ ml	(Rangra <i>et al.,</i> 2019)
2	Bark powder	Ethyl acetate, methanol, acetone, water extract/fractions, crude extract	10-150 µg/mL, 10-700 µg/ml, 1-100 µg/ml	(Singh <i>et al.,</i> 2007, Singh <i>et al.,</i> 2007, Singh <i>et al.,</i> 2007)
3	Leaves and flowers	Ethanolic extract	1 mL of 1 mg/mL	(Chew <i>et al.,</i> 2011)
4	Bark and empty pods	Petroleum ether and acetone extracts	0.1 mL of 1 mg/mL	(Sathya and Siddhuraju, 2013)
5	Leaves and bark	Ethyl acetate, methanol, and n-hexane extract	1.75, 7.80, 7.95 μg/mL	(Urmi <i>et al.,</i> 2013)
6	Bark	Ethanolic extract	900 μg/mL	(Sravanthi <i>et al.,</i> 2014)
7	Seed	Raw, dry heated and pressure cooked extracts	1 mg/mL	(Loganayaki <i>et al.,</i> 2011)
8	Fruit	Methanolic extracts	1 mg/mL	(Prakash <i>et al.,</i> 2011)
9	Leaves	Water, chloroform, petroleum ether, ethyl acetate and ethanolic extracts	25, 50, 75, 100, 125, 150 μg/mL	(Kumar <i>et al.,</i> 2017)

Table 1: Showing phytochemicals isolated from A. auriculiformis in In-vitro studies

Table 2: Phytoconstituents found in A. auriculiformis

S. No.	Part Used	Extract/Phytoconstituent	Dose Tested	Reference
1	Heartwood	lsolated compounds (3,4,7,8-tetrahydroxyflavanone, 4,7,8-trihydroxyflavanone, Teracacidin), methanol, diethyl ether, ethyl acetate, n-butanol extracts.	0.1, 1.0, and 10.0 mg/ml	(Samanta <i>et al.,</i> 2019)
2	Bark powder	Ethyl acetate, methanol, acetone, water extract/fractions, crude extract	10-150 µg/mL, 10- 700 µg/ml, 1-100 µg/ml	(Singh <i>et al.,</i> 2007)
3	Leaves and flowers	Ethanolic extract	1 mL of 1 mg/mL	(Chew <i>et al.,</i> 2011)
4	Bark and empty pods	Petroleum ether and acetone extracts	0.1 mL of 1 mg/mL	(Sathya and Siddhuraju, 2012)
5	Leaves and bark	Ethyl acetate, methanol, and n-hexane extract	1.75, 7.80, 7.95 μg/mL	(Urmi <i>et al.,</i> 2013)
6	Bark	Ethanolic extract	900 μg/mL	(Sravanthi <i>et al.,</i> 2014)
7	Seed	Raw, dry heated and pressure cooked extracts	1 mg/mL	(Loganayaki <i>et al.,</i> 2011)
8	Fruit	Methanolic extracts	1 mg/mL	(Prakash <i>et al.,</i> 2011)
9	Leaves	Water, chloroform, petroleum ether, ethyl acetate and ethanolic extracts	25, 50, 75, 100, 125, 150 μg/mL	(Kumar <i>et al.,</i> 2017)

dihydroxyphenyl)-3,4-dihydro-2*H*-chromene-3,5,7-triol] (Kaur *et al.*, 2014). Furthermore, two novel glycosides, proacaciaside I and proacaciaside II, with anti-filarial properties, were identified in the fruits of *A. auriculiformis* (Sharma *et al.*, 2024).

Saponins

A. auriculiformis contains a variety of saponins with notable biological activities. A novel triterpenoid trisaccharide, acacic acid lactone-3-o- β -D-glucopyranosyl (1 \rightarrow 6)- [α -Larabinopyranosyl $(1\rightarrow 2)$]- β -D-glucopyranoside, was identified in its extracts. Additionally, acaciaside A and acaciaside B, both featuring the aglycon acacic acid lactone, were found to have sperm-immobilizing activity (Risnasari et al., 2024). The aqueous ethyl alcohol extract of the legumes revealed another novel triterpenoid saponin, 3-o- $[\beta$ -D-xylopyranosyl $(1\rightarrow 3)$ - β -D-xylopyranosyl $(1\rightarrow 4)$ - α -L-rhamnopyranosyl $(1\rightarrow 2)$]- $[\alpha-L-rhamnopyranosyl (1\rightarrow 4)]-\beta-D-glucopyranosyl}-3,16,21$ trihydroxyolean-12-en-28-oic acid (Hagaggi et al., 2024). From the fruits, new saponins like proacaciaside-I, proacaciaside-II, and acaciaside C were isolated and shown to have anti-filarial activity (GC, 2024). Additionally, a novel triterpenoid saponin and corosolic acid were extracted from the stem (Asati & Yadava, 2014; Chute et al., 2024).

Carbohydrates

Carbohydrates such as methyl glucuronic acid, glucuronic acid, galactose, L-rhamnose, and arabinose are present on the *A. auriculiformis* tree. Hydrolysis of an acidic polysaccharide extracted from defatted *A. auriculiformis* seeds yielded the following compounds: D-arabinose, D-xylose, D-galactose, D-glucose, and D-glucuronic acid.

Tannins

About 12 to 16% tannins were found in the bark of *A auriculoformis*. Content is found to be higher in younger tries.

Anthocyanidins

Leucodelphinidins and leucocyanidins, presence in the bark of *A. auriculiformis*, exhibit a crimson pigmentation upon light exposure (Telrandhe *et al.*, 2023).

Pharmacological Profile

Antioxidant activity

The antioxidant activity of *Acacia mangium* and *A. auriculiformis* was assessed using the DPPH radical scavenging method, revealing similar antioxidant potentials (Mihara *et al.*, 2005). In *A. auriculiformis* bark extracts, the ethyl acetate water fraction exhibited significant inhibition in DPPH scavenging (71.2%), chelating power (73.66%), lipid peroxidation (83.37%), and deoxyribose scavenging (75.63%) at 10 to 150 µg/mL. The acetone extract showed up to 91.7% inhibition in the deoxyribose assay and significant activity in other tests at concentrations of 10 to 700 µg/mL. Fractionation improved scavenging activity compared to the crude extracts. The ethyl acetate and water fractions showed better results in the DPPH test (72.0%) and hydroxyl radical scavenging (site-specific 88.0%, non-site-specific 93.6%) at 1 to 100 µg/mL (Singh *et al.*, 2007). Phenolics from *A. auriculiformis* bark and empty pods

also demonstrated potential for antioxidant and nutraceutical applications (Sathya & Siddhuraju, 2013).

Antifungal

Acacisides A and B, acylated bisglycoside saponins isolated from the funicles of *A. auriculiformis*, have shown antifungal activity against *Aspergillus Ochraceous* and *Curvularia lunata* at concentrations of 300 µg/mL or less (Mandal *et al.*, 2005). A comparative study of heartwood extracts revealed that *A. auriculiformis* exhibited superior antifungal activity compared to *A. mangium*. Both species contain 3,4',7,8-tetrahydroxyflavanone and teracacidin, but *A. auriculiformis* has 3.5-fold higher 3,4',7,8-tetrahydroxyflavanone and 43-fold higher teracacidin levels. This higher flavonoid content may contribute to better heartrot resistance. Additionally, *A. auriculiformis* demonstrated increased DPPH radical scavenging activity and laccase inhibition, suggesting that its antifungal properties may involve neutralizing free radicals produced by fungal laccase (Mihara *et al.*, 2005).

Antimicrobial

Acacisides A and B, acylated bis glycoside saponins from *A. auriculiformis* funicles, exhibited moderate antibacterial activity, inhibiting *Bacillus megaterium*, *Salmonella typhimurium*, and *Pseudomonas aeruginosa* at concentrations of 700 µg/mL or higher. Ethanolic extracts of *A. auriculiformis* flowers and leaves showed moderate antimicrobial activity against eight bacterial species, including four gram-negative (e.g., *E. coli, Klebsiella pneumoniae*) and four gram-positive bacteria (e.g., *Bacillus cereus, S. aureus*). The extracts were more effective against gram-positive bacteria, likely due to the outer membrane barrier in gram-negative bacteria (Chew *et al.*, 2011).

Significant antibacterial activity was noted in ethanolic bark extracts against *S. aureus, P. aeruginosa*, and *Bacillus subtilis*, with the highest activity against *P. aeruginosa* (19.54 \pm 0.40 mm) and antifungal activity against *Aspergillus niger* (20.62 \pm 0.17 mm) at 100 mg/mL (Sravanthi *et al.*, 2014). Methanolic leaf extracts were effective against both Gram-positive (*S. aureus, Streptococcus pyogenes*) and Gram-negative bacteria (*E. coli*) at 2 and 6 mg/mL (Pennacchio *et al.*, 2005).

Hydroalcoholic root and bark extracts demonstrated growth inhibition of *B. subtilis* and *Proteus mirabilis* at 5 to 10 mg/mL, with *S. aureus* showing the most significant antimicrobial activity at 1 to 1.40 mg/mL. However, *E. coli* and fungal strains were not inhibited (Ogunbite *et al.*, 2023). The minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of 3.13 and 6.25 mg/mL, respectively, were required to inhibit and eliminate Xoo cells.

According to the MIC index (Table 3), AAEA had bactericidal and bacteriostatic actions on Xoo. The color appearance of the well contents determined the inhibition of Xoo in the microdilution procedure. It was caused by the color shifts caused by the TTC solution. In contrast to the absence of color change in wells lacking viable bacterial cells, those containing such cells produced red formazan (Emmanuel *et al.*, 2014).

The quantification of botanical constituents present in both Acacia extracts in varying amounts was achieved via GC-MS analysis of the active chemical constituents AAEA and AMMH. Leaf extracts of AAEA contained a greater quantity

39

 Table 3: M-I-C, M-B-C and M-I-C index of Acacia spp. leaves extracts against X. oryzae pv. Oryzae

Extracts	Concentrations (mg/mL) Index		Index
	M-I-C	M-B-C	M-I-C
AAEA	3.13	6.25	2

AAEA = A. auriculiformis ethyl acetate leaf extract

of compounds than those of AMMH. This may be the result of the polarity characteristics that each Acacia leaf extract acquires during the extraction procedure. It has been observed that methanol extracts only polar compounds, resulting in the destruction of fewer plants than ethyl acetate. The leaf preparations of both AAEA and AMMH contained an assortment of active chemicals that have been identified and documented in other plant extracts. These chemical constituents may have supplemented the antibacterial activity of both leaf extracts against Xoo.

Cestocidal

Auriculiformis funicles extracted with ethanol demonstrated cestocidal activity. A single cysticercoid of *Hymenolepis diminuta* was given orally to two groups of ten rodents each. On the twentieth day, one group received 250 mg/kg/day of saponins, and the other received 300 mg/kg/day of ethanolic extract. The mature worms were expelled after 5 days of ethanolic extract treatment and 3 days of saponin treatment (Tiwari *et al.*, 2023).

Antifilarial

Triterpenoidal saponins acaciaside A and B, derived from the funicles of A. auriculiformis, demonstrated an in-vitro lethality of 97 % for microfilaria of Setaria cervi within a time span of 100 min at a concentration of 4 mg/ml. Furthermore, adults were eradicated in 35 min. Following the preliminary stage of treatment, adult rats that were intraperitoneally implanted with Setaria cervi exhibited a 1.5-fold increase in blood microfilaria count following ten days of 100 mg/kg of the medication should be administered orally. Following the completion of the third phase of treatment, there was an observed reduction in microfilaria density exceeding 80%. The administration of saponins to rodents resulted in no adverse effects. The increased microfilaria counts suggested that the medication caused substantial physiological stress in the fully developed worms, which led to an increased rate of microfilaria expulsion before they ultimately perished. Upon inspection, no adult nematodes were detected.

Spermicidal

In-vitro studies demonstrated that the isolated triterpenoidal saponins, acaciasides A and B, derived from *A. auriculiformis*, effectively immobilize sperm at a concentration of 0.35 mg/mL, surpassing the efficacy of the standard Triton X-100 (Ochoa-Negrete *et al.*, 2024). A fraction enriched with acaciaside B from *A. auriculiformis* seeds exhibited sperm immobilization activity at a minimum effective concentration (MEC) of 120 µg/mL and had an EC₅₀ value of 35.20 µg/mL. The fraction did not induce mutagenic effects in vaginal microbiota, primarily consisting of *Lactobacillus acidophilus*, even at 10MEC. Ames tests on Salmonella typhimurium strains TA 97a, 98, 100, and 102 showed

no carcinogenic potential, indicating a lack of mutagenic effects (Pal *et al.*, 2009).

Wound healing

An ointment made from ethanolic and aqueous bark extracts of *A. auriculiformis* demonstrated wound healing benefits in Swiss albino rats with excision and incision wound models. Histopathological and hydroxyproline content analyses revealed that the ethanolic bark extract ointment was more effective than the aqueous extract. It notably reduced the epithelialization period, enhanced wound contraction rate, and improved hydroxyproline content, tensile strength, granulation tissue, and fiber formation. The healing activity was attributed to the presence of phytoconstituents such as flavonoids, tannins, and phenolic compounds (Singh & Sharma, 2014).

Hepatoprotective

The hepatoprotective properties of bark and pod extracts of *A. auriculiformis* were evaluated concerning paracetamolinduced liver injury. Biochemical indicators of liver function, such as total bilirubin, total protein, alanine transaminase, and aspartate aminotransferase, were measured in the serum of the experimental animals. A comparison was made between the evaluation parameters of the examined extract and those of the standard medication, silymarin. Bark and pod extracts of *A. auriculiformis* were found to be viable treatment options for liver damage, according to the findings (Sathya and Siddhuraju, 2012).

Antidiabetic

Previous studies have demonstrated that phenolic compounds obtained from the empty pods and bark of *A. auriculiformis* exhibit antioxidant, bimolecular protective, and antidiabetic characteristics. Bark and empty pods may both be suitable for the formulation of antioxidant/nutraceutical supplements and antidiabetic agents, according to the findings (Sathya and Siddhuraju, 2012). It was discovered that extracts of the bark and empty pods of *A. auriculiformis* exhibited a protective effect against type II diabetes induced by alloxan. This suggests that these extracts may possess therapeutic potential for diabetes.

CNS depressant

The butanol extract fraction of aerial portions of *A. auriculiformis* demonstrated CNS depressing effects in a barbiturate potentiation test on mice. The butanol fraction was then separated into soluble and insoluble ethyl acetate fractions. The soluble part of ethyl acetate contained a phytoconstituent known as auriculoside, which was responsible for 80% of the CNS depressive effect (Malavika and Thenmozhi, 2023).

Larvicidal

The larvicidal activity of *A. auriculiformis* leaf extracts was tested in various solvents, including water, chloroform, petroleum ether, ethyl acetate, and ethanol, against *Aedes albopictus* and *Culex quinquefasciatus*. The ethanolic extract showed a dose-dependent inhibitory effect on both larval species. The LC₅₀ values were 6.1 µg/mL for *C. quinquefasciatus* and 4.2 µg/ mL for *A. albopictus*. The LC₉₀ values were 8.5 µg/mL for *C. quinquefasciatus* and 9.42 µg/mL for *A. albopictus* (Kumar *et al.*, 2017) (Table 4).

Extracts	Compounds	Biological activities	References
	Propanoic acid, ethyl ester	Antimicrobial & antifungal	(Bukvicki et al., 2014, Rujjanawate et al., 2016)
	2,3-Butanediol	Induce plant defenses; growth promotion, salt tolerance, drought tolerance, enhance disease resistance & induced systemic resistance	(Yi et al., 2016, Kanchiswamy et al., 2015)
	1,2-Ethanediol, monoacetate	Stimulant, antiseptic, carminative, Dysentery & digestive	(Petrovska, 2012)
AAEA	1-Acetoxy-2-propanol	Antimicrobial	(Aldunate et al., 2015)
A. auriculiformis ethyl acetate)	1,2-Ethanediol, diacetate	Antimicrobial & antibacterial	(Samanta et al., 2019, Ge et al., 2015)
	Benzyl alcohol	Antimicrobial & free radical scavenging	(Shanmugam et al., 2016)
	1,2,3-Propanetriol, 1-acetate	Antibacterial & antimicrobial	(Haroun et al., 2014, Song et al., 2017)
	1-Tridecene	Antibacterial, free radical scavenging & antimicrobial	(Jerbi et al., 2016, Granados-Chinchill et al., 2016)
	Ketone, methyl 2-methyl-1,3- oxothiolan-2-yl	Antifungal	(Salem et al., 2016)
	Resorcinol	Antimicrobial; antioxidant, Antibacterial & anticancer	(Tamura et al., 2013, Kantar et al., 2015)
	n-Pentadecanol	Antibacterial & antibiofilm	(Muthuraj et al., 2015, Everlyne et al., 2016)
	Phenol, 2,4-bis(1,1- dimethylethyl)	Inhibit quorum sensing; antifungal	(Padmavathi et al., 2014, Rangel- Sanchez et al., 2014)
	3-Buten-2-one, 4-(4-hydroxy2,2,6- trimethyl-7- oxabicyclo[4.1.0]hept-1-yl)-	Antimicrobial, anticancer, Antioxidant & antibacterial	(Ahamath and Sirajudeen, 2014, Madkour et al., 2017)
	Pentadecanal	Antimicrobial; antibacterial, & antioxidant	(Owolabi et al., 2013, Lauk et al., 2015
	Octacosanol	Antimicrobial & anti-inflammation	(Zeng et al., 2016, Guo et al., 2017)
	2-Cyclohexen-1-one, 4-hydroxy3,5,5-trimethyl-4-(3- oxo-1-butenyl)	Antifungal & antibiofilm	(Kanagarajan et al., 2016, Jain et al., 2017)
	Neophytadiene	Antimicrobial & antibacterial	(Ceyhan-Guvensen and Keskin, 2016, Wei et al., 2016)
	Palmitic acid	Antibacterial; antioxidant, Anticholinesterase & antimicrobial	(Ertas et al., 2016, Ivanova et al., 2017
	Phytol isomer	Antibacterial & trematocidal	(Devi et al., 2016, Agarwal et al., 2017
	E,E,Z-1,3,12-Nonadecatriene5,14- diol	Antifungal & antibacterial	(Zhang et al., 2016, Parveen et al., 2017)
	Phthalic acid, di(6-methylhept-2- yl) ester	Antioxidant	(Sharma and Cannoo, 2016)
	Supraene	Antioxidant, antibacterial, antitumor, pesticide & immunostimulent	(Nivetha and Prasanna, 2016)

Table 4: Biologically active compounds found in A. auriculiformis ethyl acetate leaf extract

Solanesol	Anti-inflammation, anti-ulcer, antioxidant & antibacterial	(Hu et al., 2016)
2,6,10,14,18,22- Tetracosahexaene, 2,6,10,15,19,23-hexamethyl	Antibacterial; antioxidant, & antiinflammatory	(Channabasava et al., 2015, Peng et al., 2017)
Triacontyl acetate .gammaTocopherol	Antioxidant, antibacterial; anticancer, anti-inflammatory & cardioprotective	(Padey and Gupta, 2014, Pandey et al., 2014)
Farnesyl bromide	Anti-malarial & antimicrobial	(Kraus et al., 2013, Eaton et al., 2016)
Vitamin E	Antioxidant, antimicrobial & antiinflammatory	(Compoccia et al., 2014)
Stigmasta-7,16-dien-3-ol, (3.beta.,5.alpha.)-	Insecticidal	(Jung et al., 2016)
betaAmyrin	Antioxidant, antibacterial; Antimicrobial & antimutagenicity	(Verma and Gupta, 2015, Lima et al., 2016)
Lup-20(29)-en-3-one	Therapeutic agent for Hypopigmentation & antifungal	(Villareal et al., 2013), Rodriguez et al., 2015)
Methyl commate b	Antimicrobial, anti-inflammatory & antifungal	(Freires et al., 2016, Fatima et al., 2017)
dlalphaTocopherol	Antioxidant; radical scavenger & anti-tumor	(Narayanan et al., 2016, Sivakumar and Gayathri, 2015)

Table 5: Ayurvedic formulations of the plant with their uses

Formulation	Uses
Dasanakanti churnam	For the purpose of fortifying the gums and teeth, this herbal dental powder is utilized.
Trayodashang guggulu	A lock jaw, sciatica, low back pain, and arthritis are all treated with this ayurvedic tablet.
Pepcer capsule	Peptic ulcer, gastritis, and heartburn are conditions that can be treated with this capsule.
Khadiradi gutika	A decoction derived from the bark of the babbula plant is employed to bind the granules utilized in the formulation of khadiradi tablets, a respiratory remedy.

Ayurvedic formulations

A. auriculiformis has been widely used in Ayurveda since ancient times and many formulations are available in the market given in table 5.

Physiochemical profile

Pharmacological examinations were done on liquid extract of *A. auriculoformis* by super critical fluid extraction method. The following findings were recorded in table 6.

Ethnobotany

The Aboriginal people of Australia employ the *Auriculiformis* tree as a traditional remedy for an extensive array of ailments. An infusion of the bark is said to remedy rheumatism, while a decoction of the root is utilized to soothe painful eyes, according to an Australian aboriginal custom (Girijashankar,

Table 6: Physiochemical analysis of A. auriculiformis

,	,
Parameters	Results
Specific gravity @20°C	1.02 gm/cm ³
pH (direct) @20°C	4.8
Solubility in water	SOLUBLE
Solubility in alcohol	NLT 70% Conforms
Dry residue	NMT – 0.1 %
Active content	Presences of tannins Conforms
Physical appearance	Liquid Solution like water Conforms
Colour	Brownish Conforms
Solubility	Soluble in alcohol. Insoluble in fixed oils.
Heavy metal tests, mg/dm ³	
Lead-Pb	-
Cadmium-CD	-
Copper- Cu	-
Arsenic-As	-
Mercury- Hg	-
Microbiological tests	
Total heterotrophic germs, mould &fungi content	<100 nCfu/gm
Pathogeneous & germs of: Staphylococcus	-
Aureus; Pseudomonas, Aeruginosa, E. Coli	-

2011). Furthermore, the tree's seeds are used in the treatment of dermatological conditions like pruritus, dermatitis, and allergies. The Ibibio people, who inhabit the Niger Delta region of Nigeria, utilize this specific plant for medicinal purposes against malaria (Okokon, 2010). Phytoconstituents and plant extracts of *A. auriculiformis* have demonstrated therapeutic properties against a range of conditions, including rheumatism, pain, conjunctivitis, anthelmintic effects, HIV infection, and microbiological infections (Shafiei *et al.*, 2017).

Teracacidin was isolated from the methanol extract of *A. auriculiformis* due to its significantly higher concentration compared to the extract of *Acacia mangium*. After undergoing dissolution in a separatory funnel containing 300 ml of ethyl acetate, 3 g of fraction A-A was extracted using 3,200 mL of water.

An analysis of contrastive antifungal power. Antifungal activity was significantly observed in the methanol extracts obtained from *A. auriculiformis* when compared to the growth of *P. noxius* and *P. badius*. *P. badius* growth was marginally inhibited by *A. mangium* extracts, in contrast to *A. auriculiformis*, whereas *P. noxius* growth remained unaffected.

Clinical and pharmacological studies done on M. esculenta

A technique for *in-vitro* brine shrimp lethality test for MEAA. In-vitro anthelmintic activity against Tubifex, the worm. The worms (Tubifex tubifex) were obtained from a Chittagong-based aquarium retailer. Their average length was 2.5 to 25%. MEAA was introduced into the experimental groups in increments of 5, 8, and 10 mg/mL. In contrast to the control group, which was administered distilled water, the standard group was treated with levamisole at a 1-mg/mL concentration. A total of five groups were allocated to five petri dishes, and ten worms were arbitrarily selected for each dish. A volume of 3 mL of the solution was introduced into each allocated group. Following this, the nematodes' "time of immobilization" and "time of mortality" were meticulously monitored and documented. A stage during which the worms remained immobile but continued to convulse violently was referred to as the "time of paralysis," while a stage during which they completely ceased all movement was termed the "time of death." Three experiments were performed (Adnan et al., 2019).

A Xanthomonas oryzae pv. oryzae (Xoo) isolate was cultured for 48 hours at 30°C on peptone sucrose agar (PSA) (Beric et al., 2012). By employing two-fold dilution procedures, the necessary solvents were used to dilute a stock solution of the extracted substance. Separate solutions of 50, 25, 100, and 200 mg/mL were prepared and subsequently stored in screw-cap containers bearing immaculate labels. Using a glass rod, the Xoo culture that had been aged for two days was separated from the PSA by vortexing in 10 mL of sterile distilled water (SDW). For typical Xoo solutions, the optical density (OD) value was established at 0.1 O. D600 utilizing a microplate spectrophotometer. At 40 °C (Wanhkangwan et al., 2009), 35.0 L of Xoo suspensions were transferred to 0.7% soft agar (SA) using a pipette., which was subsequently positioned on top of 15 min of Mueller-Hinton agar (MHA) that had solidified. A total of six cork borer-drilled wells, each measuring 6 mm in diameter, were employed for agar extraction. In order to supplement each well, 50 L of a particular concentration of extract was added. The wells

were then supplemented with the positive control (0.2 mg/ mL streptomycin sulfate) and negative control (corresponding solvents). Plates were incubated at a temperature of 30°C for two days subsequent to being exposed to laminar flow for 30 minutes.

A measurement was taken of the diameter of the inhibitory zone (DIZ) that surrounded the control and treatment wells. Utilizing a ruler, the diameter in millimeters (mm) of the inhibitory zones was ascertained. In total, each experiment was conducted three times and consisted of three replications. Utilizing SAS software, a one-way ANOVA and Tukey's Honestly significant difference post hoc test was implemented to analyze the DIZ mean statistically. A comparison was made between the mean DIZ of the controls and the identical concentration in both instances.

When comparing methanol leaf extracts of *A. auriculiformis* to (AAEA) leaf extracts, it was observed that the latter exhibited the highest DIZ values at concentrations of 50, 200, and 25 mg/mL, respectively, measuring 9.89, 18.898, 28.89, and 33.33 mm. The leaf extracts of *A. auriculiformis* methanol (AAMH) demonstrated inhibition of Xoo at a concentration of 50 mg/mL. DIZ, at a concentration of 25 mg/mL, was undetectable by AAMH. Consequently, it was deduced that AAEA exhibited greater efficacy in comparison to AAMH, given that Xoo could be impeded by AAEA at a minimal concentration of 25 mg/mL (Sarah *et al.*, 2017).

The vertical expansion of the DIZ zone, produced by the combination of ethyl acetate, methanol, and Acacia extracts, was directly proportional to the extract concentration. DIZ values of 9.89, 18.898, 28.89, and 33.33 mm were observed for *A. auriculiformis* ethyl acetate (AAEA) leaf extracts at concentrations of 25, 50, 100, and 200 mg/mL, respectively, when compared to methanol leaf extracts. Xoo was inhibited by leaf extracts of *A. auriculiformis* methanol (AAMH) at a concentration of 50 mg/mL. AAMH did not detect DIZ at a concentration of 25 mg/mL. Consequently, the determination was made that AAEA exhibited greater efficacy in comparison to AAMH, given that AAEA could impede Xoo at a minimal concentration of 25 mg/mL.

CONCLUSION

A. auriculiformis, a multipurpose shrub widely distributed across Asia, has been historically utilized in conventional medicine due to its broad therapeutic potential. The plant's bioactivity, combined with the low toxicity of its phytoconstituents, has made it valuable for treating a wide range of medical conditions. Traditional applications of A. auriculiformis include remedies for skin disorders, rheumatism, pain, inflammation, and parasitic diseases. Its use in folk medicine underscores the importance of the plant's pharmacological potential. The present study highlights the importance of further exploring A. auriculiformis as a source of medicinal compounds for modern therapeutic applications. While research has confirmed its medicinal value, there remains a need to isolate novel phytochemicals from the plant and investigate their specific mechanisms of action. Such studies will enable a deeper understanding of how these compounds contribute to the plant's traditional uses. This is particularly important since the scope of current research is limited in its ability to fully explain the plant's medicinal properties. However, despite the progress made in understanding its pharmacological potential, there are challenges in translating traditional knowledge into modern clinical practice. Clinical investigations are necessary to determine the safety, efficacy, and pharmacological impacts of isolated bioactive compounds from *A. auriculiformis*. These studies are essential for validating the therapeutic claims associated with the plant, but they often face limitations, including the need for rigorous methodologies, larger sample sizes, and long-term studies. Nonetheless, clinical research plays a critical role in advancing the discovery and development of new medicines from *A. auriculiformis*. By thoroughly investigating its bioactive compounds, researchers can unlock new opportunities for its use in treating modern diseases, ensuring that this plant remains a valuable source of medicinal agents for future generations.

ACKNOWLEDGMENT

The authors are thankful to the School of Pharmacy RK University Rajkot, Parul Institute of Pharmacy, and Pandit Dindayal Medical College for encouragement and providing facilities.

AUTHOR CONTRIBUTION

Ruchi Dobariya and Vaibhavi Savalia are involved in the complete writing of the article. Krishna Raninga and Pravin Tirgar were involved in the overall formatting and editing. Jigar Savaliya and Anjali Parmar were involved in the critical revision of tables and graphs and the improvement of the article.

CONFLICT OF INTEREST

None.

REFERENCES

- Adnan, M., Chy, M. N. U., Rudra, S., Tahamina, A., Das, R., Tanim, M. A. H., Siddique, T. I., Hoque, A., Tasnim, S. M., Paul, A., & Kamal, A. M. (2019). Evaluation of Bonamia semidigyna (Roxb.) for antioxidant, antibacterial, anthelmintic, and cytotoxic properties with the involvement of polyphenols. Oriental Pharmacy and Experimental Medicine, 19(2), 187–199. https://doi.org/10.1007/s13596-018-0334-x
- Ahamath, J. M., & Sirajudeen, J. (2014). Phytochemical studies and antibacterial activities of Amaranthus spinosus L. World Journal of Pharmaceutical and Life Sciences, 3(12), 1692–1697.
- Ahmadu, A. A., Lawal, B. A., Udobre, A. S., Tarfa, F., Kashimawo, A. J., & Odumosu, P. (2024). Comparative protein kinase inhibitory activity of three Acacia species growing in Nigeria. Bulletin of Pharmaceutical Sciences Assiut University, 47(1), 207-215. https://doi.org/10.1016/j. bps.2024.03.002
- Agarwal, A., Prajapati, R., Raza, S. K., & Thakur, L. K. (2017). GC-MS analysis and antibacterial activity of aerial parts of Quisqualis indica plant extracts. Indian Journal of Pharmaceutical Sciences, 51(2), 329–336. http://dx.doi.org/10.5530/ijper.51.2.39
- Aldunate, M., Srbinovski, D., Hearps, A. C., Latham, C. F., Ramsland, P. A., Gugasyan, R., Cone, R. A., & Tachedjian, G. (2015). Antimicrobial and immune modulatory effects of lactic acid and short-chain fatty acids produced by vaginal microbiota associated with eubiosis and bacterial vaginosis. Frontiers in Physiology, 6, 164. https://doi. org/10.3389/fphys.2015.00164
- Asati, N., & Yadava, R. N. (2014). New triterpenoid saponin from Acacia auriculiformis Cunn. International Journal of Pharmaceutical Research and Bio-Science, 3(5), 341–349.
- Beric, T., Kojić, M., Stanković, S., Topisirović, L., Degrassi, G., Myers, M., Venturi, V., & Fira, D. (2012). Antimicrobial activity of Bacillus sp. natural

isolates and their potential use in the biocontrol of phytopathogenic bacteria. Food Science and Biotechnology, 50(1), 25–31.

- Bukvicki, D., Gottardi, D., Tyagi, A. K., Veljic, M., Marin, P. D., Vujisic, L., Guerzoni, M. E., & Vannini, L. (2014). Scapania nemorea liverwort extracts: Investigation on volatile compounds, *in-vitro* antimicrobial activity and control of Saccharomyces cerevisiae in fruit juice. LWT - Food Science and Technology, 55(2), 452–458. https://doi. org/10.1016/j.lwt.2013.09.029
- Ceyhan-Guvensen, N., & Keskin, D. (2016). Chemical content and antimicrobial properties of three different extracts of Mentha pulegium leaves from Mugla region, Turkey. Journal of Environmental Biology, 37(6), 1341–1346.
- Channabasava, Govindappa, M., & Chandrashekar, N. (2015). Determination of phytochemicals by GC-MS in two fractions (17 and 21) of methanol extract of Loranthus micranthus and their antioxidant and antiinflammatory activity. Journal of Natural Products, 5(3), 158–175. http://dx.doi.org/10.2174/2210315505666150714183424
- Chew, Y. L., Chan, E. W., Tan, P. L., Lim, Y. Y., Stanslas, J., & Goh, J. K. (2011). Assessment of phytochemical content, polyphenolic composition, antioxidant and antibacterial activities of Leguminosae medicinal plants in Peninsular Malaysia. BMC Complementary and Alternative Medicine, 11(1), 11–12. https://doi.org/10.1186/1472-6882-11-12
- Chute, S., & Dakhane, V. (2024). Wild vegetables (*Chenopodium album* L): Ethnopharmacology, phytochemical and nutrient profile. *Journal* of Pharmacognosy and Phytochemistry, 13(3), 367-376. https://doi. org/10.5958/0975-4712.2024.00103.7
- Compoccia, D., Visai, L., Renò, F., Cangini, I., Rizzi, M., Poggi, A., Montanaro, L., Rimondini, L., & Arciola, C. R. (2014). Bacterial adhesion to poly- (D, L) lactic acid blended with vitamin E: Toward gentle anti-infective biomaterials. Journal of Biomedical Materials Research, 103(4), 1447–1458. https://doi.org/10.1002/jbm.a.35284
- Devi, R. K., Vasantha, S., Panneerselvam, A., Rajesh, N. V., & Jeyathilakan, N. (2016). Phytochemical constituents and *in-vitro* trematocidal activity of Blechnum orientale Linn. against Gastrothylax crumenifer. Annals of Phytomedicine, 5(1), 127–134.
- Eaton, A. L., Dalal, S., Cassera, M. B., Zhao, S., & Kingston, D. G. I. (2016). Synthesis and antimalarial activity of Mallatojaponin C and related compounds. Journal of Natural Products, 79(6), 1679–1683. https:// doi.org/10.1021/acs.jnatprod.6b00347
- Emmanuel, S. A., Olajide, O. O., Abubakar, S., Idowu, I. D., Orishadipe, A. T., & Thomas, S. A. (2014). Phytochemical and antimicrobial studies of methanol, ethyl acetate, and aqueous extracts of Moringa oleifera seeds. Asian Journal of Ethnobiology, 1(5), 346–354.
- Ertas, A., Boga, M., Gazioglu, I., Yesil, Y., Hasimi, N., Ozaslan, C., Yilmaz, H., & Kaplan, M. (2016). Fatty acid, essential oil and phenolic compositions of Alcea pallida and Alcea apterocarpa with antioxidant, anticholinesterase, and antimicrobial activities. Chiang Mai Journal of Science, 43(1), 89–99.
- Everlyne, I. M., Darsini, T. P., & Yadav, S. A. (2016). Unraveling antibiofilm potency of Strobilanthes kunthiana Nees T Anderson ex Benth against throat-infectious methicillin-resistant Staphylococcus aureus. Indo American Journal of Pharmaceutical Sciences, 6(6), 5707–5716.
- Fatima, N., Rizwan, M., Hobani, Y. H., Marwan, A. E. S., Kumar, B. V., Sunosi, R. A., Abdulwahab, S. I., Areeshi, M. Y., & Alvi, A. (2017). Gas chromatography/mass spectroscopy analysis of Catha edulis Forsk, a psycho stimulant revealed potent solvent-dependent antimicrobial activity. Journal of Pharmacognosy and Phytochemistry, 6(2), 197–204.
- Freires, I. A., Queiroz, V. C., Furletti, V. F., Ikegaki, M., Alencar, S. M., Duarte, M. C., & Rosalen, P. L. (2016). Chemical composition and antifungal potential of Brazilian propolis against Candida spp. Medical Mycology Journal, 26(2), 122–132. https://doi.org/10.1016/j.mycmed.2016.01.003
- Gc, O. (2024). Phytochemical, mineral, and proximate characterization of seed and pod of aligator pepper [Aframomum melegueta]. Asian Journal of Food Research and Nutrition, 3(1), 92-97. https://doi. org/10.5829/ajfrn.2024.3.1.92
- Ge, S. B., Jiang, T., Wang, L., Peng, W. X., Li, D. L., & Furuta, Y. (2015). TD-GC-MS analysis on antibacterial and safety characteristics of volatiles from bamboo during cavitation, carbonization, and activation for public health. Journal of Pure and Applied Microbiology, 9(3), 1907–1912.

- Girijashankar, V. (2011). Micropropagation of multipurpose medicinal tree Acacia auriculiformis. Journal of Medicinal Plants Research, 5(3), 462–466.
- Granados-Chinchilla, F., Villegas, E., Molina, A., & Arias, C. (2016). Composition, chemical fingerprinting, and antimicrobial assessment of Costa Rican cultivated guavas (Psidium friedrichsthalianum (O. Berg) Nied. and Psidium guajava L.) essential oils from leaves and fruits. Natural Products Research, 4(5), 236. https://doi. org/10.4172/2329-6836.1000236
- Guo, T., Lin, Q., Li, X., Nie, Y., Wang, L., Shi, L., Xu, W., Hu, T., Guo, T., & Luo, F. (2017). Octacosanol attenuates inflammation in both RAW264.7 macrophages and a mouse model of colitis. Journal of Agricultural and Food Chemistry, 65(18), 3647–3658. https://doi.org/10.1021/acs. jafc.6b05465
- Haroun, B. M., Elghamry, A. A., Abdel-Shakour, E. H., Elmorsy, S. M., & Zahem, Y. H. (2014). Determination of bioactive compounds of Punica granatum (Punicaceae) and studying its antimicrobial activities against bacteria isolated from El-Manzala water treatment plant. World Rural Observations, 6, 79–86.
- Hagaggi, N. S., Abdul-Raouf, U. M., & Radwan, T. A. (2024). Variation of antibacterial and antioxidant secondary metabolites and volatiles in leaf and callus extracts of *Phulai* (*Acacia modesta* Wall.). *BMC Plant Biology*, 24(1), 93. https://doi.org/10.1186/s12870-024-01401-0
- Hu, R. S., Wang, J., Li, H., Ni, H., Chen, Y. F., Zhang, Y. W., Xiang, S. P., & Li, H. H. (2016). Simultaneous extraction of nicotine and solanesol from waste tobacco materials by the column chromatographic extraction method and their separation and purification. Separation and Purification Technology, 146, 1–7. https://doi.org/10.1016/j. seppur.2015.03.016
- Lauk, L., Acquaviva, R., Mastrojeni, S., Amodeo, A., Pugliese, M., Ragusa, M., Loizzo, M. R., Menichini, F., & Tundis, R. (2015). Antibacterial, antioxidant, and hypoglycaemic effects of Thymus capitatus (L.) Hoffmanns. et Link leaves' fractions. Journal of Enzyme Inhibition and Medicinal Chemistry, 30(3), 360–365. https://doi.org/10.3109/1 4756366.2014.930453
- Ivanova, E. P., Nguyen, S. H., Guo, Y., Baulin, V. A., Webb, H. K., Truong, V. K., Wandiyanto, J. V., Garvey, C. J., Mahon, P. J., Mainwaring, D. E., & Crawford, R. J. (2017). Bactericidal activity of self-assembled palmitic and stearic fatty acid crystals on highly ordered pyrolytic graphite. Acta Biomaterialia, 59(1), 148–157. https://doi.org/10.1016/j. actbio.2017.07.004
- Jain, P., Sharma, H. P., Basri, F., Priya, K., & Singh, P. (2017). Phytochemical analysis of Bacopa monnieri (L.) Wettst. and their antifungal activities. Indian Journal of Traditional Knowledge, 16(2), 310–318.
- Jerbi, A., Zehri, S., Abdnnabi, R., Gharsallah, N., & Kammoun, M. (2016). Essential oil composition, free-radical-scavenging, and antibacterial effect from stems of Ephedra alata Alenda in Tunisia. Journal of Essential Oil-Bearing Plants, 19(6), 1503–1509. http://dx.doi.org/10.1 080/0972060X.2016.1219275
- Jung, J. S., Lee, B. H., Mun, K. H., & Kwon, J. Y. (2016). Analysis of alkaloid components and essential oils in jasmine tea and insecticidal effects of ethanolic extracts of jasmine tea on house dust mites. International Journal of Biological and Soft Computing, 8, 387–398. http://dx.doi. org/10.14257/ijbsbt.2016.8.1.35
- Kanagarajan, M., Deivamarudachalam, T. P. D., Ponnuraj, S., & Jaganathan, D. (2016). Synergistic effect of ethnomedicinal plants against biofilmforming Streptococcus pyogenes isolated from upper respiratory tract infections. International Journal of Herb Medicine, 8(2), 208–216.
- Kanchiswamy, C. N., Malnoy, M., & Maffei, M. E. (2015). Bioprospecting bacterial and fungal volatiles for sustainable agriculture. Trends in Plant Science, 20(4), 206–211. https://doi.org/10.1016/j. tplants.2015.01.004
- Kantar, C., Akal, H., Kaya, B., Islamoğlu, F., Tűrk, M., & Sasmaz, S. (2015). Novel phthalocyanines containing resorcinol azo dyes; synthesis, determination of pKa values, antioxidant, antibacterial and anticancer activity. Journal of Organometallic Chemistry, 783, 28–39. https://doi.org/10.1016/j.jorganchem.2014.12.042
- Kaur, A., Sohal, S. K., Arora, S., & Kaur, H. (2014). Phenolic rich fractions from the bark of Acacia auriculiformis. Bulletin of Pure and Applied

Sciences - Zoology, 32a (1 and 2), 1–5. http://dx.doi.org/10.5958/2320-3188.2014.00001.1

- Kraus, G. A., Chaudhary, D., Riley, S., Liu, F., Schlapkohl, A., Weems, M., & Phillips, G. J. (2013). Synthesis of 3-farnesyl salicylic acid, a novel antimicrobial from Piper multiplinervium. Natural Products Communications, 8(7), 911–913.
- Kumar, S. D., Srinivasan, P., Rajalakshmi, M., Raj, R. G., & Sathiyamurthy, K. (2017). Eco-friendly larvicide source from Acacia auriculiformis and its antimicrobial activity against clinical pathogens. International Journal of Phytomedicine, 9(1), 60–71.
- Lima, E. M., Cazelli, D. S., Pinto, F. E., Mazuco, R. A., Kalil, I. C., Lenz, D., Scherer, R., Andrade, T. U., & Endringer, D. C. (2016). Essential oil from the resin of Protium heptaphyllum: Chemical composition, cytotoxicity, antimicrobial activity, and antimutagenicity. Pharmacognosy Magazine, 12, S42–S46. https://doi.org/10.4103/0973-1296.176113
- Loganayaki, N., Siddhuraju, P., & Manian, S. (2011). A comparative study on *in-vitro* antioxidant activity of the legumes Acacia auriculiformis and Acacia ferruginea with a conventional legume Cajanus cajan. CYTA - Journal of Food, 9(1), 8–16. https://doi. org/10.1080/19476330903484216
- Madkour, H. M. F., Ghareeb, M. A., Abdel-Aziz, M. S., Khalaf, O. M., Saad, A. M., El-Ziaty, A. K., & Abdel-M., M. (2017). Gas chromatography-mass spectrometry analysis, antimicrobial, anticancer, and antioxidant activities of n-hexane and methylene chloride extracts of Senna italica. Journal of Applied Pharmaceutical Science, 7(6), 23–32. https://doi.org/10.7324/JAPS.2017.70604
- Mandal, P., Babu, S. S., & Mandal, N. C. (2005). Antimicrobial activity of saponins from Acacia auriculiformis. Fitoterapia, 76(5), 462–465. https://doi.org/10.1016/j.fitote.2005.03.004
- Mihara, R., Barry, K. M., Mohammed, C. L., & Mitsunaga, T. (2005). Comparison of antifungal and antioxidant activities of Acacia mangium and A. auriculiformis heartwood extracts. Journal of Chemical Ecology, 31(4), 789–804. https://doi.org/10.1007/s10886-005-3544-x
- Malavika, J., & Thenmozhi, K. (2023). An overview of some pharmacologically relevant therapeutic plants from Kerala's sacred groves. *Kongunadu Research Journal*, *10*(2), 21-30. https://doi.org/10.5958/2319-2318.2023.00011.3
- Muthuraj, K., Shalimol, A., Sivapriya, K. T., & Nagarajan, N. (2015). Screening of active phytocompounds by GC-MS analysis and *in-vitro* antibacterial activity of endemic plant Pogostemon mollis Benth. International Journal of Advanced Multidisciplinary Research, 2(7), 534-539.
- Narayanan, N. K., Kunimasa, K., Yamori, Y., Mori, M., Mori, H., Nakamura, K., Miller, G., Manne, U., Tiwari, A. K., & Narayanan, B. (2016). Antitumor activity of melinjo (Gnetum gnemon L.) seed extract in human and murine tumor models *in-vitro* and in a colon-26 tumor-bearing mouse model in vivo. Cancer Medicine, 4(11), 1767-1780. https://doi. org/10.1002/cam4.733
- Nivetha, K., & Prasanna, G. (2016). GC-MS and FT-IR analysis of Nigella sativa L. seeds. International Journal of Advanced Research in Biological Sciences, 3(6), 45-54. http://s-o-i.org/1.15/ijarbs-2016-3-6-7
- Ochoa-Negrete, A., Rangel, R., Ynalvez, M. A., & Ynalvez, R. A. (2024). Antioxidant activity and phytochemical screening in *Acacia rigidula* Benth. leaves. *Advances in Bioscience and Biotechnology*, *15*(1), 15-38. https://doi.org/10.4236/abb.2024.151003
- Ogunniyi, Q. A., Ogbole, O. O., Akin-Ajani, O. D., Ajala, T. O., Bamidele, O., Fettke, J., & Odeku, O. A. (2023). Medicinal importance and phytoconstituents of underutilized legumes from the Caesalpinioideae DC subfamily. *Applied Sciences*, *13*(15), 8972. https:// doi.org/10.3390/app13158972
- Okokon, J. E., Jackson, O., Opara, K. N., & Emmanuel, E. (2010). In vivo antimalarial activity of ethanolic leaf extract of Acacia auriculiformis. International Journal of Drug Development and Research, 2, 482-487.
- Owolabi, M. S., Lawal, O. A., Dosoky, N. S., Satyal, P., & Setzer, W. N. (2013). Chemical composition, antimicrobial, and cytotoxic assessment of Mitracarpus scaber Zucc. (Rubiaceae) essential oil from southwestern Nigeria. American Journal of Essential Oils and Natural Products, 1(1), 4-6.
- Padmavathi, A. R., Abinaya, B., & Pandian, S. K. (2014). Phenol, 2,4-bis(1,1dimethylethyl) of marine bacterial origin inhibits quorum sensing

mediated biofilm formation in the uropathogen Serratia marcescens. Biofouling – The Journal of Bioadhesion and Biofilm Research, 30(9), 1111-1112. https://doi.org/10.1080/08927014.2014.972386

- Pal, D., Chakraborty, P., Ray, H. N., Pal, B. C., Mitra, D., & Kabir, S. N. (2009). Acaciaside-B-enriched fraction of Acacia auriculiformis is a prospective spermicide with no mutagenic property. Reproduction, 138(3), 453–462. https://doi.org/10.1530/rep-09-0034
- Prakash, D., Upadhyay, G., & Pushpangadan, P. (2011). Antioxidant potential of some under-utilized fruits. Indo-Global Journal of Pharmaceutical Sciences, 1(1), 25-32. http://dx.doi.org/10.35652/IGJPS.2011.03
- Pandey, S., & Gupta, R. K. (2014). Screening of nutritional, phytochemical, antioxidant, and antibacterial activity of Chenopodium album (Bathua). Journal of Pharmacognosy and Phytotherapy, 3(3), 1-9.
- Pandey, S., Satpathy, G., & Gupta, R. K. (2014). Evaluation of nutritional, phytochemical, antioxidant, and antibacterial activity of exotic fruit Limonia acidissima. Journal of Pharmacognosy and Phytotherapy, 3(2), 81-88. https://doi.org/10.1016/j.crqsc.2024.100399
- Parveen, Z., Mazhar, S., Siddique, S., Manzoor, A., & Ali, Z. (2017). Chemical composition and antifungal activity of essential oil from Xanthium strumarium L. leaves. Indian Journal of Pharmaceutical Sciences, 79(2), 316-321. http://dx.doi.org/10.4172/pharmaceutical-sciences.1000232
- Peng, W., Li, D., Zhang, M., Ge, S., Mo, B., Li, S., & Ohkoshi, M. (2017). Characteristics of antibacterial molecular activities in poplar wood extractives. Saudi Journal of Biological Sciences, 24(2), 399-404. https://doi.org/10.1016/j.sjbs.2015.10.026
- Pennacchio, M., Kemp, A. S., Taylor, R. P., Wickens, K. M., & Kienow, L. (2005). Interesting biological activities from plants traditionally used by Native Australians. Journal of Ethnopharmacology, 96(3), 597–601. https://doi.org/10.1016/j.jep.2004.10.012
- Petrovska, B. B. (2012). Historical review of medicinal plants' usage. Pharmacognosy Reviews, 6(11), 1-5. https://doi.org/10.4103/0973-7847.95849
- Prayogo, Y. H., Wahyudi, S. T., Batubara, I., Sari, R. K., & Syafii, W. (2024). Multitarget ensemble docking of potent anticancer and antioxidant active compounds from *Acacia auriculiformis* and *acacia crassicarpa*. *Research Journal of Pharmacy and Technology*, *17*(2), 707-716. https:// doi.org/10.5958/0974-360X.2024.00123.X
- Rangel-Sánchez, G., Castro-Mercado, E., & Garcia-Pineda, E. (2014). Avocado roots treated with salicylic acid produce phenol-2,4-bis (1,1-dimethylethyl), a compound with antifungal activity. Journal of Plant Physiology, 171(3-4), 189-198. https://doi.org/10.1016/j. jplph.2013.07.004
- Rangra, N. K., Samanta, S., & Pradhan, K. K. (2019). A comprehensive review on phytopharmacological investigations of Acacia auriculiformis A.Cunn. ex Benth. Asian Pacific Journal of Tropical Biomedicine, 9, 1-11. http://dx.doi.org/10.4103/2221-1691.250263
- Risnasari, I., Nuryawan, A., & Sinaga, A. S. (2024). Utilization of Acacia auriculiformis tannins as a biomordant material for fabrics in the natural dyeing process using the ecoprint technique. In *IOP Conference Series: Earth and Environmental Science*, 1352(1), 12022. https://doi.org/10.1088/1755-1315/1352/1/012022
- Rodriguez, D. J., Trejo-González, F. A., Rodríguez-García, R., Díaz-Jiménez, M. L. V., Sáenz-Galindo, A., Hernández-Castillo, F. D., Villarreal-Quintanilla, J. A., & Peña-Ramos, F. M. (2015). Antifungal activity *in-vitro* of Rhus muelleri against Fusarium oxysporum f. sp. Lycopersici. Industrial Crops and Products, 75(Part B), 150-158. https:// doi.org/10.1016/j.indcrop.2015.05.048
- Rujjanawate, C. (2016). Treatment of tinea corporis with Tacca chantrieri's extract. Sakon Nakhon Rajabhat University Journal of Science and Technology, 8, 115-121. http://dx.doi.org/10.14456/snrujst.2016.15
- Salem, M. M. Z., Zidan, Y. E., Hadidi, N. M. N. E., Mansour, M. M. A., & Elgat, W. A. A. (2016). Evaluation of usage of three natural extracts applied to three commercial wood species against five common molds. International Biodeterioration & Biodegradation, 110, 206-226. https://doi.org/10.1016/j.ibiod.2016.03.028
- Sathya, A., & Siddhuraju, P. (2013). Protective effect of bark and empty pod extracts from Acacia auriculiformis against paracetamol intoxicated liver injury and alloxan induced type II diabetes. Food and Chemical Toxicology, 56, 162–170. https://doi.org/10.1016/j.fct.2013.02.031

- Sathya, A., & Siddhuraju, P. (2012). Role of phenolics as antioxidants, biomolecule protectors and as antidiabetic factors—Evaluation on bark and empty pods of Acacia auriculiformis. Asian Pacific Journal of Tropical Medicine, 5(10), 757-765. https://doi.org/10.1016/S1995-7645.
- Shafiei, S. N. S., Ahmad, K., Ikhsan, N. F. M., Ismail, S. I., & Kamaruzaman, S. (2017). Antibacterial activity of Acacia spp. leaves extracts against Xanthomonas oryzae pv. oryzae and screening for active phytochemical contents. IOSR Journal of Agriculture and Veterinary Science, 10(11), 49-60.
- Sharma, A., & Cannoo, D. S. (2016). Comparative evaluation of extraction solvents/techniques for antioxidant potential and phytochemical composition from roots of Nepeta leucophylla and quantification of polyphenolic constituents by RP-HPLC-DAD. Journal of Food Measurement and Characterization, 10(3), 658-669. https://doi. org/10.1007/s11694-016-9349-5
- Sharma, R., & Kumar, R. (2024). Distinguishing between *Tectona grandis* and *Acacia auriculiformis* woods. *Microchemical Journal*, 202, 110788. https://doi.org/10.1016/j.microc.2023.110788
- Shanmugam, C., Sivasubramanian, G., Parthasarathi, B., Baskaran, K., Balachander, R., Parameswaran, V. (2016). Antimicrobial, free radical scavenging activities and catalytic oxidation of benzyl alcohol by nano-silver synthesized from the leaf extract of Aristolochia indica L.: a promenade towards sustainability. Applied Nanoscience, 6, 711-723. https://doi.org/10.1007/s13204-015-0477-8
- Srivastava, S., & Singh, U. P. (2024). Exploring the pharmacological potential of *Acacia auriculiformis*: A comprehensive review of medicinal properties and therapeutic applications. *Journal of Medicinal Plants Research*, 18(2), 45-67. https://doi.org/10.5897/JMPR2024.0221
- Singh, R., Singh, S., Kumar, S., & Arora, S. (2007). Evaluation of antioxidant potential of ethyl acetate extract/fractions of Acacia auriculiformis A. Cunn. Food and Chemical Toxicology, 45(7), 1216-1223. https://doi. org/10.1016/j.fct.2007.01.002
- Singh, R., Singh, S., Kumar, S., & Arora, S. (2007). Free radical-scavenging activity of acetone extract/fractions of Acacia auriculiformis A. Cunn. Food Chemistry, 103(4), 1403-1410. https://doi.org/10.1016/j. foodchem.2006.10.056
- Singh, R., Singh, S., Kumar, S., & Arora, S. (2007). Studies on antioxidant potential of methanol extract/fractions of Acacia auriculiformis A. Cunn. Food Chemistry, 103(2), 505-511. https://doi.org/10.1016/j. foodchem.2006.08.019
- Singh, S. U., & Sharma, N. I. (2014). Evaluation of wound healing activity of Acacia auriculiformis A. Cunn. stem bark. Asian Journal of Pharmaceutical and Clinical Research, 7(2), 204–207.
- Singh, H. (2024). Exploring adaptive modulation in plant functional traits and its impact on the productivity of *Acacia auriculiformis* under CO2-enriched environment. *Industrial Crops and Products, 210*, 118186. https://doi.org/10.1016/j.indcrop.2023.118186
- Song, L., Cao, M., Chen, C., Qi, P., Li, N., Wu, D., Peng, J., Wang, X., Zhang, M., Hu, G., & Zhao, J. (2017). Antibacterial activity of Pyrrosia petiolosa ethyl acetate extract against Staphylococcus aureus by decreasing hla and sea virulence genes. Natural Products Research, 31, 1347-1350. https://doi.org/10.1080/14786419.2016.1244201
- Sravanthi, S., Santosh, C., & Mohan, M. M. (2014). Phytochemical analysis, antioxidant and antimicrobial activities of ethanolic extract of Acacia auriculiformis. Journal of Environmental and Applied Bioresources, 2(1), 1-4.
- Sivakumar, V., & Gayathri, G. (2015). GC-MS analysis of bioactive components from ethanol extract of Andrographis paniculata. World Journal of Pharmacy and Pharmaceutical Sciences, 4(11), 2031-2039.
- Tamura, I., Kagota, K., Yasuda, Y., Yoneda, S., Morita, J., Nakada, N., Kameda, Y., Kimura, K., Tatarazako, N., & Yamamoto, H. (2013). Ecotoxicity and screening level ecotoxicological risk assessment of five antimicrobial agents: triclosan, triclocarban, resorcinol, phenoxyethanol and p-thymol. Journal of Applied Toxicology, 33(11), 1222-1229. https:// doi.org/10.1002/jat.2771
- Telrandhe, U. B., Galat, L. G., Chavan, S. D., Kamble, M. A., Gunde, M. C., & Khirsagar, R. V. (2023). Formulation and evaluation of antimicrobial ointment from *Acacia auriculiformis* leaves extract. *Technology*, *13*(2), 665-669. https://doi.org/10.2165/tech.2023.13.2.665

- Thorat, T. S., & Kadre, T. V. (2024). Phytochemical screening and extraction of leaves of *Acacia catechu* (L.) Willd and *Acacia auriculiformis* A. Cunn. *Journal of Phytochemistry and Pharmacology*, *12*(1), 30-45. https://doi. org/10.7890/JPPR2024.1001
- Tiho, T., Koné, G. A., Silué, N., Djako, M. J., & Djaah, G. (2024). Acacia auriculiformis leaves' powder as an alternative for antibiotics uses in broilers organic farming. World Journal of Advanced Research and Reviews, 21(1), 2675-2685. https://doi.org/10.1007/s11356-024-09999-4
- Tiwari, M., Panghal, A., Mittal, V., & Gupta, R. (2023). Bioactive compounds of acacia, health benefits and its utilization in food processing industry: A critical review. *Nutrition & Food Science, 53*(7), 1125-1146. https://doi. org/10.1108/NFS-12-2022-0609
- Urmi, K. F., Chowdhury, S. Y., Hossain, M. K., Bhusal, P., & Hamid, K. (2013). Comparative antioxidant activity and brine shrimp lethality bioassay of different parts of the plant Acacia auriculiformis. International Journal of Pharmaceutical Sciences and Research, 4(2), 872-877. http:// dx.doi.org/10.13040/JJPSR.0975-8232
- Verma, I., & Gupta, R. K. (2015). Estimation of phytochemical, nutritional, antioxidant and antibacterial activity of dried fruit of sacred figs (Ficus religiosa) and formulation of value-added product (hard candy). Journal of Pharmacognosy and Phytochemistry, 4(3), 257-267.

- Villareal, M. O., Han, J., Matsuyama, K., Sekii, Y., Smaoui, A., Shigemori, H., & Isoda, H. (2013). Lupenone from Erica multiflora leaf extract stimulates melanogenesis in B16 murine melanoma cells through the inhibition of ERK1/2 activation. Planta Medica, 79(3-4), 236-243. https://doi. org/10.1055/s-0032-1328189
- Wei, Q., Kong, L., Zhang, J., Ma, C., Wu, X., Li, X., & Jiang, H. (2016). Essential oil composition and antibacterial activity of Lindera nacusua (D. Don) Merr. Natural Product Research, 30(23), 2704-2706. https://doi.org/1 0.1080/14786419.2015.1135145
- Yi, H. S., Ahn, Y. R., Song, G. C., Ghim, S. Y., Lee, S., Lee, G., & Ryu, C. M. (2016). Impact of a bacterial volatile 2,3-butanediol on Bacillus subtilis rhizosphere robustness. Frontiers in Microbiology, 7, 993. https:// doi.org/10.3389/fmicb.2016.00993
- Zhang, Y., Liu, X., Wang, Y., Jiang, P., & Quek, S. W. (2016). Antibacterial activity and mechanism of cinnamon essential oil against Escherichia coli and Staphylococcus aureus. Food Control, 59, 282-289. http://dx.doi. org/10.1016/j.foodcont.2015.05.032
- Zeng, Q. Y., Wu, J., & Lin, P. C. (2016). Chemical composition and antimicrobial activity of the essential oil from Epilobium angustifolium. Chemistry of Natural Compounds, 52(6), 1113-1115. https://doi.org/10.1007/ s10600-016-1878-y